

GEO THERMALISM IN THE PROVINCE OF VERONA

Lake Garda and Verona Province case studies

Laura Agostini¹, Collareda M.³, Castellaccio E.⁴, Dal Degan D.⁴, Antonio Galgaro¹⁻², Jacopo Boaga¹

¹Department of Geosciences – University of Padua, Padua (Italy) laura.agostini2@studenti.unipd.it

² Institute for Geosciences and Earth Resources, IGG-CNR Padua, Italy

³⁻⁴Geologists of Vicenza (Italy) and Verona

1. Introduction

The presence of territories characterized by local temperature and heat flux anomalies, as the Southern Garda Lake and the Verona Province areas, is well known (Fig.1a;1b). They are found in the Po Plain, located in northern Italy between the Alpine and Apennine mountain ranges. The use of low and medium temperature geothermal resources enhances a sustainable development of the area, combined with significant energy and cost savings, as well as a substantial reduction of greenhouse emissions into the atmosphere. To promote a proper use of geothermal energy through direct uses of heat and by coupling with heat pump systems it is essential to characterize the geothermal reservoirs involved, taking into account the local differences related to geological, hydrogeological and thermal features.




Fig. 2 – The hydrogeological background model of district A modified (Sorbini et al., 1984).




Fig.1a –Map of temperatures and permeable rocks in the Po Plain with main hydrothermal systems. (Pasquale, Verdoya and Chiozzi, 2014)
In evidence Verona's Province (Fig.1 b)




Fig.1b – Thermal district map of the Verona Province (Castellaccio and Zorzin 2012)

2.Tectonic and Structural Characteristic

The relationship between tectonic - structural characteristic, and the presence of thermal fluids is not clear and for this reason we are studying:

- the difference between the chemical-physical compositions of thermal groundwater of the two districts identified from literature (districts A and B in Fig.1b);
- the presence of thermal contamination within the Lake Garda;
- the hydrothermal circuits for assessing the geothermal resource renewability.
- In the past authors such as Sighinolfi and Sorbini (Fig. 2) helped produce geochemical and hydrogeological models that are still worthy of interest. The hydrogeological profile in the figure represents the thermal area of Caldiero called District A in Castellaccio e Zorzin, 2012. (fig.2)

3. Geochemical and Geological data

The groundwater geochemical characterization obtained from the screening of available literature (Castellaccio & Zorzin 2012; Sighinolfi et al., 1982) shows the presence of different geochemical facies. The Piper diagram of the districts highlights the existence of calcium and magnesium bicarbonate waters differentiated by their temperature level ranging between 24 and 31 °C for Caldiero groundwaters and 12 and 14°C for those belonging to the Lessinian mountains (Fig.3a). In the B district instead (Fig.3b), thermal waters, are divided into four groups according to their different temperature. In August 2013 Further analysis were carried out (put in evidence in Piper diagram, Fig.3a and 3b). At the end of september new isotopic geochemical data will be analyzed to complement previous studies.




Fig.3a – Chemical analysis of wells and springs district A. In evidence the close correlation between analysis of Lessinian mountains cold waters and warm waters 2013 data




Fig.3b – Analysis of district B. Thermal waters with different temperatures are highlighted. The chemical analysis of 2013 are marked in green (Castellaccio and Zorzin, 2012 modified by L.Agostini)

4.Methodology

Seismic Sounding and Remote Sensing, passive seismic survey as HVSR (Horizontal Vertical Spectral Ratio), analysis of satellite and terrestrial images LiDar) will be performed to improve the local knowledge of the geothermal resources.




Fig.4a Tromograph and tromographic device for HVSR acquisition





Fig.4b Tromographic data in geological map. In evidence the Rocca hill and the Gazzo hill, whose origin is volcanic.

Fig.4c Two examples of HVSR output of Giunone spa in Caldiero town. We can see a high peak when rock is compact and a large peak when rock is fractured such as near area volcanic (the Rocca hill and the Rocca hill in Caldiero- see Fig.4b)



Fig.5 Image of Eastern Verona Province (LiDar Ministero dell'Ambiente e della Tutela del Territorio e del Mare) modified by A. Ninfo –Geosciences-UniPd

5. Further developments

In the research programme, some thermal analyses images have been performed to define the thermal characterization of the districts and a detailed investigation about the characterization of Caldiero thermal area with a tromograph (see Fig.4b). Terrestrial thermography, seismic sounding and new geochemical analyses will be carried out in the lake area and the Verona Province to better define the hydrothermal circuit of the thermal areas and to foresee a rational and sustainable use of the resources.

6. References

- Antonelli R. e Stefanini S., 1983 - Nuovi contributi idrogeologici ed idrochimici sugli acquiferi dell'Alta Pianura Veronese. Mem. Sci. Geol., 35 (1982-83), pp.35-67, figg.22.
- Castellaccio E. e Zorzin R. a cura di, 2012 - Acque calde e geotermia della provincia di Verona. Aspetti geologici e applicazioni- Memorie del Museo Civico di Storia Naturale di Verona, 2. serie - Sezione Scienze della Terra n. 8, anno 2012
- Dal Degan D., 2000 - Aspetti geologici, idrogeologici e gestionali sull'idrotermalismo nei complessi alluvionali e fluvio-glaciali Nord-Ovest della città di Verona. Alcuni Confronti con altri campi idrotermali del Veneto. Tesi di laurea inedita, a.a. 1999-2000, Università di Padova
- Gambillara R., Astori A., Massironi M., Martin S., 2003-Neotettonica dell'area veronese: uno studio integrato fra microrilievo e remote sensing- Mem. Sc. Geol. di Padova Vol.55, pp. 13-20. ISSN: 0391-8602
- Patrizi G., 2002 – Sintesi dei risultati dell'indagine idrogeologica, geochemica e geochemica-isotopica sugli acquiferi della Lessinia. Relazione inedita, Servin –Servizi Integrati Gestionali Ambientali, Ravenna
- Piccoli G., Bellati R., Binotti C., Di Lallo E., Sede R., Dal Pra' A., Cataldo R., Gatto G.O., Ghezzi G., Marchetti M., Bulgarelli G., Schiesaro G., Panichi C., Tongiorgi E., Baldi P., Ferrara G.C., Massari F., Medizza F., Illicito V., Norinelli A., DE VECCHI Gp., Gregnanin A., Piccirillo E.M., Sbettega G., 1977 - Il sistema idrotermale euganeo-berico e la geologia dei Colli Euganei. Mem. Istituti Geol. Mineral. Univ. Padova, 30 (1975 - 77), pp. 1-266, figg. 97, tabb. 14., tav. 1,1 c. geol. color. (1:25.000)
- Sighinolfi G.P., Gorgoni C., Martinelli G., Sorbini L., 1982. Indagine geochemica preliminare sulle acque del sistema termale veronese. Energia geotermica, CNR, Progr. Fin. Energia, 3: 13-20, Roma
- Thank to all authors of book " Acque calde e geotermia della provincia di Verona. Aspetti geologici e applicazioni" : Agostini Laura – laura.agostini2@studenti.unipd.it, agoalb@tiscali.it; Castellaccio Enrico – ecastellaccio@tiscali.it; Collareda Matteo – info@studiosisma.com; Dal Degan Davide – davidedaldegan@alice.it; Galadini Fabrizio – galadini@ingv.it; Micheletti Luca – artemi.progettazioni@libero.it; Monegato Giovanni – g.monegato@csg.to.cnr.it; Rogledi Sergio – sergio.rogledi@eni.com; Scardia Giancarlo – giancarlo.scardia@igag.cnr.it; Zorzin Roberto – roberto.zorzin@comune.verona.it; Minciotti Giuseppe (Director of Museo of Natural History of Verona) – giuseppe.minciotti@comune.verona.it